

IDM2008 - Identification of Dark Matter 2008

Contribution ID : 376

Measurement of Dark Matter Content at the LHC (15'+5')

Monday 18 Aug 2008 at 16:35 (00h20')

Recent cosmological observations have introduced tremendous constraint on supersymmetry models. The minimal supergravity (mSUGRA) model explains the observed dark matter content of the universe and requires stau-neutralino co-annihilation mechanism in the early universe for a large region of supersymmetry parameter space. This co-annihilation region is characterized by a small mass difference ($\sim 5-15$ GeV) between the lightest stau and the lightest neutralino. An accurate measurement of the small mass difference at a collider is crucial to confirm the co-annihilation mechanism. We show that from measurements at the LHC it is possible to confirm this mechanism and using these measurements one can predict the dark matter relic density with an uncertainty of 6% with 30 fb⁻¹ of data. This is possible by introducing measurements involving b-quark jets to determine the SUGRA parameters A_0 and $\tan(\beta)$. We measure all the mSUGRA parameters with high accuracy and our methods provide precision mass measurements of the gauginos, squark, and lighter stau without the mSUGRA assumption.

This talk is based on the following papers: arXiv:0802.2968 (to appear in Phys Rev Lett); Phys.Lett.B649:73-82,2007, Phys.Lett.B639:46-53,2006.

Primary authors : Dr. DUTTA, Bhaskar Duta (Texas A&M University)

Co-authors :

Presenter : Dr. DUTTA, Bhaskar Duta (Texas A&M University)

Session classification : Connections with accelerators

Track classification : Connections with accelerators

Type : Talk